Structures of not-finitely graded Lie algebras related to generalized Heisenberg-Virasoro algebras *

Guangzhe Fan*, Chenhong Zhou*, Xiaoqing Yue*,†
* Department of Mathematics, Tongji University, Shanghai 200092, China
E-mail: yzfanguangzhe@126.com, zhouchenhong66@163.com, xiaoqingyue@tongji.edu.cn

arXiv:1409.6282v5 [math.RA] 16 Jul 2016

Abstract: In this paper, we study the structure theory of a class of not-finitely graded Lie algebras related to generalized Heisenberg-Virasoro algebras. In particular, the derivation algebras, the automorphism groups and the second cohomology groups of these Lie algebras are determined.

Key words: not-finitely graded Lie algebras, generalized Heisenberg-Virasoro algebras, derivations, automorphisms, 2-cocycles.

Mathematics Subject Classification (2010): 17B05, 17B40, 17B65, 17B68.

1 Introduction

The Heisenberg-Virasoro algebra contains the classical Heisenberg algebra and Virasoro algebra as subalgebras. As the universal central extension of the Lie algebra of differential operators on a circle of order at most one, the Heisenberg-Virasoro algebra is an important object in mathematics and physics, and has been widely studied. For example, the twisted Heisenberg-Virasoro algebra was firstly studied by E. Arbarello et al. in [1]. Various generalizations of the Heisenberg-Virasoro algebra have been extensively studied by several authors (e.g., [2, 8–10]). However, it seems to us that little has been known on not-finitely graded aspect of generalized Heisenberg-Virasoro algebras.

In recent years, some researches on Lie algebras concerning their derivation algebras, automorphisms, second cohomology groups have been undertaken by many authors (see, e.g., [3–5, 11–17]). It is well known that central extensions, which are determined by second cohomology groups, are closely related to the structures of Lie algebras (see, e.g., [6, 7]). The computation of the second cohomology groups seems to be important and interesting. Not-finitely graded Lie algebras are important objects in Lie theory, whose structure and representation theories are subjects of studies with more challenges than those of finitely graded Lie algebras. In [3], the authors have studied the structure theory of a class of not-finitely graded Lie algebras related to generalized Virasoro algebras.

In this paper, we consider the following Lie algebras, which are referred to as *generalized Heisenberg-Virasoro algebra* $HV(\Gamma)$: Let Γ be any nontrivial additive subgroup of $\mathbb{C}$, and $\mathbb{C}[\Gamma \times \mathbb{Z}_+]$ the semigroup algebra of $\Gamma \times \mathbb{Z}_+$ with basis $\{x^{\alpha,i} := x^\alpha t^i \,|\, \alpha \in \Gamma, i \in \mathbb{Z}_+\}$ and product $x^{\alpha,i}x^{\beta,j} = x^{\alpha+\beta,i+j}$. Let ∂_x, ∂_t be the derivations of $\mathbb{C}[\Gamma \times \mathbb{Z}_+]$ defined by $\partial_x(x^{\alpha,i}) = \alpha x^{\alpha,i}$, $\partial_t(x^{\alpha,i}) = ix^{\alpha,i-1}$ for $\alpha \in \Gamma$, $i \in \mathbb{Z}_+$. Denote $\partial = \partial_x + \partial_t$. Then $HV(\Gamma)$ is the Lie algebra with the underlying space $\mathbb{C}[\Gamma \times \mathbb{Z}_+]\partial \oplus \mathbb{C}[\Gamma \times \mathbb{Z}_+]$, basis $\{L_{\alpha,i} := x^{\alpha,i}\partial,$

* Supported by the National Natural Science Foundation of China (No. 11371278, 11271284) and the grant no. 12XD1405000 of Shanghai Municipal Science and Technology Commission.

† Corresponding author: X. Yue (xiaoqingyue@tongji.edu.cn).

$H_{\beta,j} := x^{\beta,j} \,|\, \alpha, \beta \in \Gamma,\ i, j \in \mathbb{Z}_+\}$ and relations

$$[L_{\alpha,i}, L_{\beta,j}] = (\beta - \alpha)L_{\alpha+\beta,i+j} + (j - i)L_{\alpha+\beta,i+j-1}, \tag{1.1}$$

$$[L_{\alpha,i}, H_{\beta,j}] = \beta H_{\alpha+\beta,i+j} + jH_{\alpha+\beta,i+j-1}, \tag{1.2}$$

$$[H_{\alpha,i}, H_{\beta,j}] = 0. \tag{1.3}$$

Let W be $\mathbb{C}[\Gamma \times \mathbb{Z}_+]\partial$ with basis $\{L_{\alpha,i} := x^{\alpha,i}\partial \,|\, \alpha \in \Gamma,\ i \in \mathbb{Z}_+\}$ and relation (1.1), and H be $\mathbb{C}[\Gamma \times \mathbb{Z}_+]$ with basis $\{H_{\beta,j} := x^{\beta,j} \,|\, \beta \in \Gamma,\ j \in \mathbb{Z}_+\}$ and relation (1.3). We simply denote $HV = HV(\Gamma)$, then W and H are subalgebras of HV. The algebra W is the generalized Witt algebra $\mathcal{W} = W(0, 1, 0; \Gamma)$ of Witt type studied in [13].

Let us recall some basic concepts. A Lie algebra $\mathcal{L}$ is *finitely graded* if there exists an abelian group G such that $\mathcal{L} = \oplus_{a\in G}\mathcal{L}_{[a]}$ is G-graded satisfying

$$[\mathcal{L}_{[a]}, \mathcal{L}_{[b]}] \subset \mathcal{L}_{[a+b]} \ \text{ and } \ \dim \mathcal{L}_{[a]} < \infty \ \text{ for } a, b \in G. \tag{1.4}$$

An element $x \in \mathcal{L}$ is ad-*locally finite* if for any $y \in \mathcal{L}$ the subspace $\mathrm{span}\{\mathrm{ad}_x^i(y) \,|\, i \in \mathbb{Z}_+\}$ is finite-dimensional, where $\mathrm{ad}_x : y \mapsto [x, y]$ $(y \in \mathcal{L})$ is the *adjoint operator* of x.

Similar to the Lie algebras studied in [3], the Lie algebra HV has the following significant features:

(1) It has a *finitely graded filtration* in the sense that there exists a filtration $0 \subset HV^{(0)} \subset HV^{(1)} \subset \cdots$, satisfying $[HV^{(i)}, HV^{(j)}] \subseteq HV^{(i+j)}$ for all $i, j \in \mathbb{Z}_+$ and each $HV^{(i)}$ is finitely graded, i.e., there exists some abelian group G which is independent of i (one can simply choose $G = \Gamma$ in this case) such that $HV^{(i)} = \oplus_{g\in G}HV_g^{(i)}$, $[HV_g^{(i)}, HV_h^{(j)}] \subseteq HV_{g+h}^{(i+j)}$, $\dim HV_g^{(i)} < \infty$ for all $g, h \in G$, $i, j \in \mathbb{Z}_+$ (cf. (1.4)) and $HV_0^{(0)} \neq 0$.

(2) It has the set $\mathfrak{F}$ of ad-locally finite elements, where

$$\mathfrak{F} = \{aL_{0,0} + \sum_{\alpha,j} b_{\alpha,j}H_{\alpha,j} \ (\text{finite sum}) \,|\, a, b_{\alpha,j} \in \mathbb{C},\ \alpha \in \Gamma, j \in \mathbb{Z}_+\}. \tag{1.5}$$

(3) HV has exactly two proper ideals: $\mathbb{C}H_{0,0}$ and H. In particular, H is the unique maximal ideal of HV (cf. Theorem 2.1).

In this paper we shall mainly study the structure theory of HV (namely, derivations, automorphisms, 2-cocycles). The Lie algebra HV is Γ-graded

$$HV = \underset{\alpha\in\Gamma}{\oplus} HV_\alpha, \quad HV_\alpha = \mathrm{span}\{L_{\alpha,i}, H_{\alpha,i} \mid i \in \mathbb{Z}_+\} \ \text{ for } \alpha \in \Gamma.$$

However, it is not finitely graded. Nevertheless, as stated in [3], due to the fact that Γ may not be finitely generated (as a group), HV may not be finitely generated as a Lie algebra. The classical techniques (such as those in [5]) cannot be directly applied to our situation

here. One must employ some new techniques in order to tackle problems associated with not-finitely graded and not-finitely generated Lie algebras (this is also one of our motivations to present our results here). For instance, one of our strategies used in the present paper is to study the actions of $L_{\alpha,0}, L_{0,i}$ respectively so that the determination of derivation algebra and automorphism group can be done much more efficiently. The main results of the present paper are summarized in Theorems 2.1, 3.1, 4.1, 4.8 and 5.1.

Throughout the paper, we denote by $\mathbb{C}$, $\mathbb{C}^*$, $\mathbb{Z}$, $\mathbb{Z}_+$, Γ^* the sets of complex numbers, nonzero complex numbers, integers, nonnegative integers, nonzero elements of Γ respectively.

2 Some properties of HV

Now we study some properties of the Lie algebra HV, which will be summarized in Theorem 2.1. Firstly, we recall the following concept.

Suppose A is a nonzero ideal of HV, and $0 \neq x = \sum_{\alpha\in\Gamma} x_\alpha \in A$ (finite sum) such that each homogeneous component $x_\alpha = \sum_{i\in\mathbb{Z}_+}(a_{\alpha,i}L_{\alpha,i} + b_{\alpha,i}H_{\alpha,i})$ (finite sum) for some $a_{\alpha,i}, b_{\alpha,i} \in \mathbb{C}$. Let

$$\mathrm{Supp}\, x = \{\alpha \in \Gamma \,|\, x_\alpha \neq 0\}, \tag{2.1}$$

then it is called the *support* of x.

Theorem 2.1. (1) *The Lie algebra HV is not finitely graded.*

(2) *HV has the set $\mathfrak{F}$ of ad-locally finite elements, where*

$$\mathfrak{F} = \{aL_{0,0} + \sum_{\alpha,j} b_{\alpha,j}H_{\alpha,j} \text{ (finite sum)} \,|\, a, b_{\alpha,j} \in \mathbb{C},\ \alpha \in \Gamma, j \in \mathbb{Z}_+\}. \tag{2.2}$$

(3) *HV has exactly two proper ideals: $\mathbb{C}H_{0,0}$ and H. In particular, H is the unique maximal ideal of HV.*

Proof (1) Firstly, we assume that there exists some abelian group G such that $HV = \oplus_{g\in G} HV_{[g]}$ is G-graded satisfying (1.4). Suppose $HV_{[0]} \neq 0$, then we can take any nonzero $x \in HV_{[0]}$ and $\beta \in \Gamma$ such that $\beta \notin \mathrm{Supp}x$ (cf. (2.1)). If we assume that $L_{\beta,2} \in \oplus_{g\in I} HV_{[g]}$ for some finite subset I of G, we obtain that $\{\mathrm{ad}_x^k L_{\beta,2} \,|\, k \in \mathbb{Z}_+\}$ is an independent subset of $\oplus_{g\in I} HV_{[g]}$. In particular, $\sum_{g\in I} \dim HV_{[g]} = \infty$. Since I is finite, there exists some $g_0 \in I$ such that $\dim HV_{[g_0]} = \infty$, contradicting our assumption that $HV_{[g]}$ is finite dimensional for any $g \in G$.

So we have $HV_{[0]} = 0$. Choose a total order "$\prec$" on G compatible with its group structure. Suppose $L_{0,0} = \sum_{i=1}^n x_{g_i}$ with $g_1 \prec \cdots \prec g_{n-1} \prec g_n$ for some $n \geq 1$, where $0 \neq x_{g_i} \in HV_{[g_i]}$. Due to

$$[L_{0,0}, L_{\alpha,0}] = \alpha L_{\alpha,0}, \tag{2.3}$$

$L_{0,0}$ preserves every one-dimensional space $\mathbb{C}L_{\alpha,0}$ ($\alpha \in \Gamma$), we have that $L_{0,0}$ can not lie in $\oplus_{0\prec g\in G}HV_{[g]}$ or $\oplus_{G\ni g\prec 0}HV_{[g]}$ of HV. So we have $0 \prec g_n$ and $g_1 \prec 0$.

By comparing the maximal homogenous components (with respect to the G-gradation of HV) of both sides of (2.3), we obtain that the maximal homogenous component of $L_{\alpha,0}$ lies in $HV_{[g_n]}$ for all $\alpha \in \Gamma$. So there exists $\lambda_\alpha \in \mathbb{C}^*$ such that $L_{\alpha,0} = \lambda_\alpha L_{0,0} + \sum_{g\prec g_n} y_g^\alpha$, where $y_g^\alpha \in HV_{[g]}$. In fact, substituting $L_{\alpha,0} = \lambda_\alpha L_{0,0} + \sum_{g\prec g_n} y_g^\alpha$ on the left hand side of (2.3) and then proceeding the analysis above gives $L_{\alpha,0} = \lambda_\alpha L_{0,0} + \sum_{g\prec 0} y_g^\alpha$. Similarly, we can deduce $L_{\alpha,0} = \lambda_\alpha L_{0,0} + \sum_{0\prec g} y_g^\alpha$. This leads to a contradiction.

(2) It can be obtained easily by the relations of HV.

(3) Let H' be a proper ideal of HV which satisfies $\mathbb{C}H_{0,0} \neq H'$, then there must exist a nonzero element $\sum_{\gamma,s} a_{\gamma,s}L_{\gamma,s} + \sum_{\beta,j} c_{\beta,j}H_{\beta,j}$ in $H' \setminus \mathbb{C}H_{0,0}$. Let any $L_{\alpha,i}$ and $H_{\alpha,i}$ act on it respectively, we get

$$\begin{aligned}[L_{\alpha,i}, \sum_{\gamma,s} a_{\gamma,s}L_{\gamma,s} + \sum_{\beta,j} c_{\beta,j}H_{\beta,j}] = & \sum_{\gamma,s} a_{\gamma,s}((\gamma-\alpha)L_{\alpha+\gamma,i+s} + (s-i)L_{\alpha+\gamma,i+s-1}) \\ & + \sum_{\beta,j} c_{\beta,j}(\beta H_{\alpha+\beta,i+j} + jH_{\alpha+\beta,i+j-1}) \end{aligned} \tag{2.4}$$

and

$$[H_{\alpha,i}, \sum_{\gamma,s} a_{\gamma,s}L_{\gamma,s} + \sum_{\beta,j} c_{\beta,j}H_{\beta,j}] = -\sum_{\gamma,s} a_{\gamma,s}(\alpha H_{\alpha+\gamma,i+s} + iH_{\alpha+\gamma,i+s-1}). \tag{2.5}$$

(i) If all the $a_{\gamma,s}$ are equal to zero, there must exist some $c_{\beta,j} \neq 0$ with $(\beta, j) \neq (0, 0)$. From (2.4), then we have $H = H'$.

(ii) If there exist some $a_{\gamma,s} \neq 0$, we can get $H' = HV$ from (2.4) and (2.5). It is a contradiction.

From the above discussion, HV has only two proper ideals: $\mathbb{C}H_{0,0}$ and H. Due to $\mathbb{C}H_{0,0} \subsetneqq H$, we get that H is the unique maximal ideal of HV. □

3 Derivation algebra of HV

Recall that a linear map $D : HV \to HV$ is a *derivation* of HV if $D\big([x, y]\big) = [D(x), y] + [x, D(y)]$ for any $x, y \in HV$. For any $z \in HV$, the adjoint operator $\mathrm{ad}_z : HV \to HV$ is a derivation, called an *inner derivation*. Denote by $\mathrm{Der}\, HV$ and $\mathrm{ad}\, HV$ the vector spaces of all derivations and inner derivations respectively. Then the first cohomology group $H^1(HV,\ HV) \cong \mathrm{Der}\, HV/\mathrm{ad}\, HV$. We say that a derivation $D \in \mathrm{Der}\, HV$ is of degree γ, if it satisfies that $D(HV_\alpha) \subset HV_{\alpha+\gamma}$. Denote by $(\mathrm{Der}\, HV)_\gamma$ the space of all derivations of degree γ.

Denote by $\mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$ the space of group homomorphisms from Γ to $\mathbb{C}$. For each $\phi \in \mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$, the scalar multiplication ϕ by $c \in \mathbb{C}$ is defined by $(c\phi)(\gamma) = c\phi(\gamma)$, thus

$\mathrm{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})$ is a vector space. Then for each $\phi\in\mathrm{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})$, we can define a derivation D_ϕ as follows:

$$D_\phi(L_{\alpha,i})=\phi(\alpha)L_{\alpha,i},\quad D_\phi(H_{\alpha,i})=\phi(\alpha)H_{\alpha,i}\quad\text{for}\ \ \alpha\in\Gamma,\ i\in\mathbb{Z}_+. \tag{3.1}$$

We still use $\mathrm{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})$ to denote the corresponding subspace of $\mathrm{Der}\,HV$. In particular, since $\phi_0:\alpha\mapsto\alpha$ is in $\mathrm{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})$, we have the derivation

$$D_0=D_{\phi_0}:L_{\alpha,i}\mapsto\alpha L_{\alpha,i},\quad H_{\alpha,i}\mapsto\alpha H_{\alpha,i}\ \ \text{for}\ \ \alpha\in\Gamma,\ i\in\mathbb{Z}_+. \tag{3.2}$$

Now we define two derivations of degree 0, which are obviously not inner derivations of HV.

$$\begin{cases} D_1(L_{\alpha,i})=\alpha H_{\alpha,i}+iH_{\alpha,i-1}, D_1(H_{\alpha,i})=0,\\ D_2(L_{\alpha,i})=0, \qquad\qquad D_2(H_{\alpha,i})=H_{\alpha,i}\ \ \text{for}\ \ \alpha\in\Gamma,\ i\in\mathbb{Z}_+.\end{cases}$$

Theorem 3.1. *The derivation space of HV can be written as*

$$\mathrm{Der}\,HV=\mathop{\oplus}_{\gamma\in\Gamma}(\mathrm{Der}\,HV)_\gamma=\mathrm{ad}HV\oplus\mathrm{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})\oplus\mathbb{C}D_1\oplus\mathbb{C}D_2, \tag{3.3}$$

where $(\mathrm{Der}\,HV)_\gamma\subset\mathrm{ad}HV$, if $\gamma\in\Gamma^*$, and $(\mathrm{Der}\,HV)_0=(\mathrm{ad}HV)_0\oplus\mathrm{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})\oplus\mathbb{C}D_1\oplus\mathbb{C}D_2$.

Proof It can be obtained by the following lemmas 3.2∼3.6 immediately. □

Lemma 3.2. *Every derivation $D\in HV$ can be written as*

$$D=\sum_{\gamma\in\Gamma}D_\gamma,\quad D_\gamma\in(\mathrm{Der}\,HV)_\gamma \tag{3.4}$$

such that only finitely many $D_\gamma(x)\neq0$ for every $x\in HV$ (such a sum in (3.4) is called summable).

Proof For a derivation $D\in\mathrm{Der}\,HV$ and an element $x_\alpha\in HV_\alpha$, assume that

$$D(x_\alpha)=\sum_{\beta\in\Gamma}y_\beta.$$

We define $D_\gamma(x_\alpha)=y_{\alpha+\gamma}$. Then a direct computation shows that D_γ is a derivation. □

Lemma 3.3. *For any $D\in\mathrm{Der}\,HV$, replacing D with $D-\mathrm{ad}_u$ for some $u\in HV$, we can suppose*

$$D(L_{0,0})=0. \tag{3.5}$$

Proof Suppose $D(L_{0,0})=\sum_{\alpha,j}(a_{\alpha,j}L_{\alpha,j}+c_{\alpha,j}H_{\alpha,j})\in HV$ for some $a_{\alpha,j},\ c_{\alpha,j}\in\mathbb{C}$. For any $\alpha\in\Gamma$, we define $b_{\alpha,j},\ d_{\alpha,j}\in\mathbb{C}$ inductively on $j\geq 0$ by

$$b_{\alpha,j}=\begin{cases} j^{-1}(-a_{\alpha,j-1}-\alpha b_{\alpha,j-1}) & \text{if } j\geq 1,\\ 0 & \text{if } j=0.\end{cases} \tag{3.6}$$

and

$$d_{\alpha,j}=\begin{cases} j^{-1}(-c_{\alpha,j-1}-\alpha d_{\alpha,j-1}) & \text{if } j\geq 1,\\ 0 & \text{if } j=0.\end{cases} \tag{3.7}$$

Take $u=\sum_{\alpha,j}(b_{\alpha,j}L_{\alpha,j}+d_{\alpha,j}H_{\alpha,j})$. Note that $u\in HV$, by (3.6) and (3.7), it gives

$$\begin{aligned} D(L_{0,0})-\mathrm{ad}_u(L_{0,0}) &= \sum_{\alpha,j}(a_{\alpha,j}L_{\alpha,j}+c_{\alpha,j}H_{\alpha,j})-\sum_{\alpha,j}(-\alpha b_{\alpha,j}-(j+1)b_{\alpha,j+1})L_{\alpha,j}\\ &\quad -\sum_{\alpha,j}(-\alpha d_{\alpha,j}-(j+1)d_{\alpha,j+1})H_{\alpha,j}\\ &= 0. \end{aligned}$$

□

Lemma 3.4. *If $\gamma\in\Gamma^*$, $D\in(\mathrm{Der}\,HV)_\gamma$ and $D(L_{0,0})=0$, then*

$$D=0. \tag{3.8}$$

Proof Applying D to $[L_{0,0},L_{\alpha,0}]=\alpha L_{\alpha,0}$ with $\alpha\in\Gamma^*$, then we have

$$[L_{0,0},D(L_{\alpha,0})]=\alpha D(L_{\alpha,0}). \tag{3.9}$$

Assume $D(L_{\alpha,0})=\sum_{j\in\mathbb{Z}_+}(b_jL_{\alpha+\gamma,j}+d_jH_{\alpha+\gamma,j})$, and by (3.9) we have

$$\begin{cases} \gamma b_j=-(j+1)b_{j+1} & \text{for } j\geq 0,\\ \gamma d_j=-(j+1)d_{j+1} & \text{for } j\geq 0.\end{cases} \tag{3.10}$$

As $b_j=0, j\gg 0$, $b_j=0$ for any $j\geq 0$. Similarly, $d_j=0$ for any $j\geq 0$. Thus we have $D(L_{\alpha,0})=0$, for any $\alpha\in\Gamma$.

Applying D to $[L_{0,0},L_{0,1}]=L_{0,0}$, then we have

$$[L_{0,0},D(L_{0,1})]=D(L_{0,0}). \tag{3.11}$$

Assume $D(L_{0,j})=\sum_{k\in\mathbb{Z}_+}(f(j,k)L_{\gamma,k}+g(j,k)H_{\gamma,k})$, for $f(j,k),\ g(j,k)\in\mathbb{C}$, and note that $f(0,k)=g(0,k)=0$. By (3.11), we have

$$\begin{cases} \gamma f(1,k)=-(k+1)f(1,k+1) & \text{for } k\geq 0,\\ \gamma g(1,k)=-(k+1)g(1,k+1) & \text{for } k\geq 0.\end{cases} \tag{3.12}$$

Then we can get $f(1,k)=g(1,k)=0$ for any $k\in\mathbb{Z}_+$, and thus $D(L_{0,1})=0$.

Now applying D to $[L_{0,0},H_{\alpha,0}]=\alpha H_{\alpha,0}$, then we have

$$[L_{0,0},D(H_{\alpha,0})]=\alpha D(H_{\alpha,0}). \tag{3.13}$$

Assume $D(H_{\alpha,0})=\sum_{j\in\mathbb{Z}_+}(e_jL_{\alpha+\gamma,j}+f_jH_{\alpha+\gamma,j})$, and from (3.13) we obtain

$$\begin{cases} \gamma e_j=-(j+1)e_{j+1} & \text{for } \ j\geq 0,\\ \gamma f_j=-(j+1)f_{j+1} & \text{for } \ j\geq 0. \end{cases} \tag{3.14}$$

As $e_j=0, j\gg 0$, $e_j=0$ for any $j\geq 0$. Similarly, $f_j=0$ for any $j\geq 0$. Thus we have $D(H_{\alpha,0})=0$, for any $\alpha\in\Gamma$.

Since HV is generated by $\{L_{\alpha,0},L_{0,1},H_{\alpha,0}\mid \alpha\in\Gamma\}$ from relation (1.1), we get $D=0$ where $D\in(\operatorname{Der}HV)_\gamma$, for $\gamma\in\Gamma^*$. □

Lemma 3.5. *Assume $D\in(\operatorname{Der}HV)_0$ and $D(L_{0,0})=0$, we have :*

(i) $D(L_{\alpha,0})=b_\alpha L_{\alpha,0}+d_\alpha H_{\alpha,0}$, where $b_\alpha\in\operatorname{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})$, $d_\alpha=\alpha d_1$, $d_1\in\mathbb{C},\alpha\in\Gamma$.

(ii) $D(H_{\alpha,0})=f_\alpha H_{\alpha,0}$, where $f_\alpha=b_\alpha+f_0$, $f_0\in\mathbb{C}$, $\alpha\in\Gamma$.

Proof For (i), assume $D(L_{\alpha,0})=\sum_{\alpha,j}(b_{\alpha,j}L_{\alpha,j}+d_{\alpha,j}H_{\alpha,j})$, and from (3.9) we have

$$\begin{cases} b_{\alpha,j}=0 & \text{for } \ j\geq 1,\\ d_{\alpha,j}=0 & \text{for } \ j\geq 1. \end{cases} \tag{3.15}$$

Simply denote $b_{\alpha,0}$, $d_{\alpha,0}$ by b_α, d_α respectively, we have $D(L_{\alpha,0})=b_\alpha L_{\alpha,0}+d_\alpha H_{\alpha,0}$, for any $\alpha\in\Gamma$. Applying D to $[L_{\alpha,0},L_{\beta,0}]=(\beta-\alpha)L_{\alpha+\beta,0}$, we get that

$$\begin{cases} b_\alpha+b_\beta=b_{\alpha+\beta} & \text{for } \ \alpha\neq\beta,\\ \beta d_\beta-\alpha d_\alpha=(\beta-\alpha)d_{\alpha+\beta} & \text{for } \ \alpha,\ \beta\in\Gamma. \end{cases} \tag{3.16}$$

As $D(L_{0,0})=b_0L_{0,0}=0$, we have $b_0=0$. According to (3.16), we also have $-b_\alpha=b_{-\alpha}$. Set $\alpha=\beta$ in (3.16), we obtain that $b_{2\alpha}=b_{(\alpha+\eta)+(\alpha-\eta)}=b_{\alpha+\eta}+b_{\alpha-\eta}=b_\alpha+b_\eta+b_\alpha+b_{-\eta}=2b_\alpha$, for any $\eta\in\Gamma\backslash\{\pm\alpha,0\}$. Thus (3.16) holds for all $\alpha,\beta\in\Gamma$, which shows that the map $\Phi:\alpha\longmapsto b_\alpha$ is an element in $\operatorname{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})$. Hence we get that $b_\alpha\in\operatorname{Hom}_{\mathbb{Z}}(\Gamma,\mathbb{C})$.

Suppose that $\Gamma'=c\Gamma$ for some $c\in\mathbb{C}^*$, we have $HV(\Gamma)\cong HV(\Gamma')$. Thus without loss of generality, we can always suppose $1\in\Gamma$. Then we can obtain that $d_\alpha=(d_1-d_0)\alpha+d_0$, for $d_1\in\mathbb{C}$. Due to $d_0=0$, we have $d_\alpha=\alpha d_1$, for $d_1\in\mathbb{C}$.

For (ii), assume $D(H_{0,0})=\sum_{j\in\mathbb{Z}_+}(e_jL_{0,j}+f_jH_{0,j})$, and apply D to $[L_{0,0},H_{0,0}]=0$. Then we have $e_j=f_j=0$, for any $j\geq 1$, and we get $D(H_{0,0})=e_0L_{0,0}+f_0H_{0,0}$. Applying D to

$[L_{0,1}, H_{0,0}] = 0$, we have $e_0 = 0$. Thus $D(H_{0,0}) = f_0 H_{0,0}$. Assume $D(H_{\alpha,0}) = \sum_{\alpha,j}(e_{\alpha,j}L_{\alpha,j} + f_{\alpha,j}H_{\alpha,j})$, and by (3.13), we have

$$\begin{cases} e_{\alpha,j} = 0 & \text{for } \ j \geq 1, \\ f_{\alpha,j} = 0 & \text{for } \ j \geq 1. \end{cases} \tag{3.17}$$

So $D(H_{\alpha,0}) = e_{\alpha,0}L_{\alpha,0} + f_{\alpha,0}H_{\alpha,0} := e_\alpha L_{\alpha,0} + f_\alpha H_{\alpha,0}$. Applying D to $[L_{-\alpha,0}, H_{\alpha,0}] = \alpha H_{0,0}$, we have $e_\alpha = 0$. Thus we can assume $D(H_{\alpha,0}) = f_\alpha H_{\alpha,0}$. Now applying D to $[L_{\alpha,0}, H_{\beta,0}] = \beta H_{\alpha+\beta,0}$, we have $b_\alpha + f_\beta = f_{\alpha+\beta}$. Therefore $b_\alpha + f_0 = f_\alpha$, with $f_0 \in \mathbb{C}$ for any $\alpha \in \Gamma$. $\square$

If we replace D' by $D - D_\phi$(cf. (3.1))(note that this replacement does not affect (3.5)), we can suppose $b_\alpha = 0$ for all $\alpha \in \Gamma$. Then we have $D'(L_{\alpha,0}) = \alpha d_1 H_{\alpha,0}$ and $D'(H_{\alpha,0}) = f_0 H_{\alpha,0}$.

Assume $D'(L_{\alpha,1}) = \sum_{j\in\mathbb{Z}_+}(f(1,j)L_{\alpha,j} + g(1,j)H_{\alpha,j})$, for $f(1,j),\ g(1,j) \in \mathbb{C}$. Applying D' to $[L_{0,0}, L_{\alpha,1}] = \alpha L_{\alpha,1} + L_{\alpha,0}$, we get $f(1,j) = g(1,j) = 0$ for $j \geq 1$. So $D'(L_{\alpha,1}) = f(1,0)L_{\alpha,0} + g(1,0)H_{\alpha,0} + \alpha d_1 H_{\alpha,1}$. By induction on i, we have $D'(L_{\alpha,i}) = if(1,0)L_{\alpha,i-1} + ig(1,0)H_{\alpha,i-1} + \alpha d_1 H_{\alpha,i}$.

Applying D' to $[L_{-\alpha,i}, H_{\alpha,0}] = \alpha H_{0,i}$ for $\alpha \in \Gamma^*$, we have $D'(H_{0,i}) = if(1,0)H_{0,i-1} + f_0 H_{0,i}$. Applying D' to $[L_{0,i}, H_{\alpha,0}] = \alpha H_{\alpha,i}$ for $\alpha \in \Gamma^*$, we obtain $D'(H_{\alpha,i}) = if(1,0)H_{\alpha,i-1} + f_0 H_{\alpha,i}$ for $\alpha \in \Gamma^*$. So $D'(H_{\alpha,i}) = if(1,0)H_{\alpha,i-1} + f_0 H_{\alpha,i}$ for $\alpha \in \Gamma,\ i \in \mathbb{Z}_+$. And it follows that

$$\begin{cases} D'(L_{\alpha,i}) = if(1,0)L_{\alpha,i-1} + ig(1,0)H_{\alpha,i-1} + \alpha d_1 H_{\alpha,i}, \\ D'(H_{\alpha,i}) = if(1,0)H_{\alpha,i-1} + f_0 H_{\alpha,i}, \end{cases}$$

for any $\alpha \in \Gamma,\ i \in \mathbb{Z}_+$. Due to $[L_{0,0}, L_{\alpha,i}] = \alpha L_{\alpha,i} + iL_{\alpha,i-1}$ and $[L_{0,0}, H_{\alpha,i}] = \alpha H_{\alpha,i} + iH_{\alpha,i-1}$, if we replace u (cf. (3.5))by $u + f(1,0)L_{0,0}$, and denote $D' + D_0$ by D' (cf. (3.2)), then

$$\begin{cases} D'(L_{\alpha,i}) = ig(1,0)H_{\alpha,i-1} + \alpha d_1 H_{\alpha,i}, \\ D'(H_{\alpha,i}) = f_0 H_{\alpha,i}, \end{cases} \tag{3.18}$$

with $g(1,0),\ f_0, d_1 \in \mathbb{C}$, for any $\alpha \in \Gamma,\ i \in \mathbb{Z}_+$.

Applying D' to $[L_{\alpha,i}, L_{\beta,j}] = (\beta - \alpha)L_{\alpha+\beta,i+j} + (j-i)L_{\alpha+\beta,i+j-1}$, we obtain $d_1 = g(1,0)$. So we have

$$\begin{cases} D'(L_{\alpha,i}) = d_1(\alpha H_{\alpha,i} + iH_{\alpha,i-1}), \\ D'(H_{\alpha,i}) = f_0 H_{\alpha,i}, \end{cases} \tag{3.19}$$

with $f_0, d_1 \in \mathbb{C}$, for any $\alpha \in \Gamma,\ i \in \mathbb{Z}_+$.

Lemma 3.6. *Under the above notations, we have a decomposition for* $(\operatorname{Der} HV)_0$:

$$(\operatorname{Der} HV)_0 = (\operatorname{ad} HV)_0 \oplus \operatorname{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}) \oplus \mathbb{C}D_1 \oplus \mathbb{C}D_2, \tag{3.20}$$

where $D_1(L_{\alpha,i}) = \alpha H_{\alpha,i} + iH_{\alpha,i-1},\ D_1(H_{\alpha,i}) = 0,\ D_2(L_{\alpha,i}) = 0,\ D_2(H_{\alpha,i}) = H_{\alpha,i}$, *for any* $\alpha \in \Gamma,\ i \in \mathbb{Z}_+$.

4 Automorphism group of HV

Firstly, we study the automorphism group of W. Denote by $\mathrm{Aut}\, W$ the automorphism group of W. Let $\chi(\Gamma)$ be the set of characters of Γ, i.e., the set of group homomorphisms $\tau : \Gamma \to \mathbb{C}^*$. Set $\Gamma^{\mathbb{C}^*} = \{c \in \mathbb{C}^* \,|\, c\Gamma = \Gamma\}$. We define a group structure on $\chi(\Gamma) \times \Gamma^{\mathbb{C}^*}$ by

$$(\tau_1, c_1) \cdot (\tau_2, c_2) = (\tau, c_1 c_2), \quad \text{where} \quad \tau : \alpha \mapsto \tau_1(c_2\alpha)\tau_2(\alpha) \text{ for } \alpha \in \Gamma. \tag{4.1}$$

It turns out that the group $\chi(\Gamma) \times \Gamma^{\mathbb{C}^*}$ is just the semidirect product $\chi(\Gamma) \rtimes \Gamma^{\mathbb{C}^*}$ under the action given by $(c\tau)(\alpha) = \tau(c\alpha)$ for all $c \in \Gamma^{\mathbb{C}^*}$, $\tau \in \chi(\Gamma)$, $\alpha \in \Gamma$. We define a group homomorphism $\phi : (\tau, c) \mapsto \phi_{\tau,c}$ from $\chi(\Gamma) \times \Gamma^{\mathbb{C}^*}$ to $\mathrm{Aut}\, W$ such that $\phi_{\tau,c}$ is the automorphism of W defined by

$$\phi_{\tau,c} : L_{\alpha,i} \mapsto \tau(\alpha) c^{i-1} L_{c\alpha,i} \quad \text{for} \quad \alpha \in \Gamma,\, i \in \mathbb{Z}_+. \tag{4.2}$$

One can easily verify that $\phi_{\tau,c}$ is indeed an automorphism of W.

Theorem 4.1. $\mathrm{Aut}\, W \cong \chi(\Gamma) \rtimes \Gamma^{\mathbb{C}^*}$.

Proof This theorem can be obtained by the following three lemmas. □

As $L_{0,0}$ is the unique locally finite element of W, for any $\sigma \in \mathrm{Aut}\, W$, we can suppose $\sigma(L_{0,0}) = c^{-1}L_{0,0}$, for $c \in \mathbb{C}^*$.

Lemma 4.2. *For any $\alpha \in \Gamma$, we have $\sigma(L_{\alpha,0}) = c^{-1}\tau(\alpha)L_{c\alpha,0}$, where $c \in \mathbb{C}^*$, $\tau \in \mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}^*)$.*

Proof Assume that $\sigma(L_{\alpha,0}) = \sum_{\beta,j} b^{\alpha}_{\beta,j} L_{\beta,j}$ (finite sum) for some $b^{\alpha}_{\beta,j} \in \mathbb{C}$. Applying σ to $[L_{0,0}, L_{\alpha,0}] = \alpha L_{\alpha,0}$, we have $[L_{0,0}, \sum_{\beta,j} b^{\alpha}_{\beta,j} L_{\beta,j}] = c\alpha \sum_{\beta,j} b^{\alpha}_{\beta,j} L_{\beta,j}$. By (1.1), we get

$$\sum_{\beta,j} b^{\alpha}_{\beta,j}(\beta L_{\beta,j} + jL_{\beta,j-1}) = c\alpha \sum_{\beta,j} b^{\alpha}_{\beta,j} L_{\beta,j}.$$

Comparing the coefficients of $L_{\beta,j}$ on both sides of the above formula, we have $\beta b^{\alpha}_{\beta,j} + (j+1)b^{\alpha}_{\beta,j+1} = c\alpha b^{\alpha}_{\beta,j}$. Then it gives that $\beta = c\alpha$, $b^{\alpha}_{\beta,j} = 0$, $j \geq 1$. So $\sigma(L_{\alpha,0}) = \varphi(\alpha)L_{c\alpha,0} = c^{-1}\tau(\alpha)L_{c\alpha,0}$, where φ is a function from Γ to $\mathbb{C}^*$ such that $\varphi(0) = c^{-1}$. Let $\tau(\alpha) = c\varphi(\alpha)$. We can verify that $\tau \in \mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C}^*)$. □

Lemma 4.3. *For any $i \in \mathbb{Z}_+$, we have $\sigma(L_{0,i}) = c^{i-1}L_{0,i}$.*

Proof The lemma holds for $i = 0$. Now we suppose it holds for $i-1$. Applying σ to $[L_{0,0}, L_{0,i}] = iL_{0,i-1}$, we have $[L_{0,0}, \sigma(L_{0,i})] = ic^{i-1}L_{0,i-1}$. Furthermore, we obtain $\sigma(L_{0,i}) = c^{i-1}L_{0,i} + aL_{0,0}$, for $a \in \mathbb{C}$. Now applying σ to $[L_{0,1}, L_{0,i}] = (i-1)L_{0,i}$ for $i > 1$, we get $a = 0$. Then it is obvious that $\sigma(L_{0,i}) = c^{i-1}L_{0,i}$. □

Lemma 4.4. *For any $\alpha \in \Gamma,\ i \in \mathbb{Z}_+$, we have $\sigma(L_{\alpha,i}) = \tau(\alpha)c^{i-1}L_{c\alpha,i}$.*

Proof If $i = 0$, the lemma holds. Now we suppose it is true for $i-1$. Applying σ to $[L_{\alpha,0}, L_{0,i}] = -\alpha L_{\alpha,i} + iL_{\alpha,i-1}$ with $\alpha \in \Gamma^*$ and using $\sigma(L_{\alpha,i-1}) = \tau(\alpha)c^{i-2}L_{c\alpha,i-1}$, we can deduce $\sigma(L_{\alpha,i}) = \tau(\alpha)c^{i-1}L_{c\alpha,i}$, for any $\alpha \in \Gamma^*,\ i \in \mathbb{Z}_+$. Then, by Lemma 4.3, we have $\sigma(L_{\alpha,i}) = \tau(\alpha)c^{i-1}L_{c\alpha,i}$, for any $\alpha \in \Gamma,\ i \in \mathbb{Z}_+$. □

Next we compute the automorphism group of H.

Lemma 4.5. *For any $\alpha \in \Gamma$, we have $\sigma(H_{\alpha,0}) = \tau(\alpha)eH_{c\alpha,0}$, where $e \in \mathbb{C}^*,\ \tau \in \mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$.*

Proof Applying σ to $[L_{0,0}, H_{\alpha,0}] = \alpha H_{\alpha,0}$, since H is an abelain ideal$(cf.(1.2), (1.3))$, we get $[L_{0,0}, \sigma(H_{\alpha,0})] = c\alpha\sigma(H_{\alpha,0})$. That is to say, $\sigma(H_{\alpha,0})$ is an eigenvector of $\mathrm{ad}_{L_{0,0}}$ with eigenvector $c\alpha$. Hence we obtain that $\sigma(H_{\alpha,0}) = f(\alpha)H_{c\alpha,0}$. Furthermore applying σ to $[L_{\alpha,0}, H_{\beta,0}] = \beta H_{\alpha+\beta,0}$ gives $\tau(\alpha)c^{-1}[L_{c\alpha,0}, f(\beta)H_{c\beta,0}] = \beta f(\alpha+\beta)H_{c(\alpha+\beta),0}$, i.e., $f(\beta)\tau(\alpha) = f(\alpha+\beta)$. Denote $f(0) = e$, we get $f(\alpha) = \tau(\alpha)e$, where $\tau \in \mathrm{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{C})$. Due to Theorem 2.1(3), then we have $e \in \mathbb{C}^*$. □

Lemma 4.6. *For any $j \in \mathbb{Z}_+$, we have $\sigma(H_{0,j}) = ec^jH_{0,j}$.*

Proof Applying σ to $[L_{0,1}, H_{0,j}] = jH_{0,j}$, and we get $[L_{0,1}, \sigma(H_{0,j})] = j\sigma(H_{0,j})$. Assume that $\sigma(H_{0,j}) = \sum_{\beta,k} x_{\beta,k}H_{\beta,k}$, we have $\sum_{\beta,k} x_{\beta,k}(\beta H_{\beta,k+1} + kH_{\beta,k}) = j\sum_{\beta,j} x_{\beta,k}H_{\beta,k}$. Comparing the coefficients of $H_{\beta,k}$ gives $\beta x_{\beta,k-1} = (j-k)x_{\beta,k}$. Thus $\beta = 0$. We can assume that $\sigma(H_{0,j}) = e_jH_{0,j}$, $e_j \in \mathbb{C}^*$. Applying σ to $[L_{0,0}, H_{0,j}] = jH_{0,j-1}$, we get $e_j = ce_{j-1},\ j > 0$. As $e_0 = e$, we have $e_j = ec^j$. Thus the lemma holds. □

Lemma 4.7. *For any $\alpha \in \Gamma, i \in \mathrm{Z}_+$, we have $\sigma(H_{\alpha,j}) = \tau(\alpha)ec^jH_{\alpha,j}$.*

Proof If $\alpha = 0$, the lemma holds. If $\alpha \neq 0$, by applying σ to $[L_{0,j}, H_{\alpha,0}] = \alpha H_{\alpha,j}$, we have $[c^{j-1}L_{0,j}, \tau(\alpha)eH_{c\beta,0}] = \alpha c^j e\tau(\alpha)H_{c\alpha,j} = \alpha\sigma(H_{\alpha,j})$. Thus we get $\sigma(H_{\alpha,j}) = \tau(\alpha)ec^jH_{\alpha,j}$. □

Now we study the automorphism group of HV. Denote by $\mathrm{Aut}\,HV$ the automorphism group of HV. Let $\sigma \in \mathrm{Aut}\,HV$. We have $\sigma(H) = H,\ \sigma(H_{0,0}) = eH_{0,0}$, for $e \in \mathbb{C}^*$. Let $\overline{\sigma} \in \mathrm{Aut}\,HV/H(i.e., \mathrm{Aut}\,W)$, and we have $\overline{\sigma(L_{\alpha,i})} = \tau(\alpha)c^{i-1}\overline{(L_{\alpha,i})}$, i.e., $\sigma(L_{\alpha,i}) \equiv \tau(\alpha)c^{i-1}L_{c\alpha,i} \bmod H$.

Since $\sigma(L_{0,0}) \equiv c^{-1}L_{0,0} \bmod H$, we can assume $\sigma(L_{0,0}) = c^{-1}L_{0,0} + \sum_{\alpha,i} a_{\alpha,i}H_{\alpha,i}$ for some $a_{\alpha,i} \in \mathbb{C}$. Define

$$b_{\alpha,i} = \begin{cases} -i^{-1}a_{\alpha,i-1} & \text{for } i > 0, \\ 0 & \text{for } i = 0. \end{cases} \tag{4.3}$$

Denote $\tau = e^{\mathrm{ad}\sum_{\alpha,i} b_{\alpha,i}H_{\alpha,i}} \in \mathrm{Inn}HV$, so $\tau(c^{-1}L_{0,0}) = c^{-1}L_{0,0} + \sum_{\alpha,i} b_{\alpha,i}H_{\alpha,i}$. Let $\pi = \tau^{-1}\sigma$. We obtain $\pi(L_{0,0}) = \tau^{-1}\sigma(L_{0,0}) = c^{-1}L_{0,0}$. Then by Lemma 4.5, 4.6 and 4.7, we have $\pi(L_{\alpha,i}) = \tau(\alpha)c^{i-1}L_{c\alpha,i}$. Thus we complete the proof of the following theorem.

Theorem 4.8. $\mathrm{Aut}\, HV \cong \mathrm{Inn} HV \rtimes \Big((\chi(\Gamma) \rtimes \Gamma^{\mathbb{C}^*}) \rtimes \mathbb{C}^*\Big)$.

5 Second cohomology group of HV

Recall that a bilinear form $\psi : HV \times HV \to \mathbb{C}$ is called a *2-cocycle* on HV if the following conditions are satisfied:

$$\psi(x,y) = -\psi(y,x), \quad \psi(x,[y,z]) + \psi(y,[z,x]) + \psi(z,[x,y]) = 0,$$

for $x, y, z \in HV$. Denote by $C^2(HV,\ \mathbb{C})$ the vector space of 2-cocycles on HV. For any linear function $f : HV \to \mathbb{C}$, one can define a 2-cocycle ψ_f by $\psi_f(x,y) = f([x,y])$ for $x, y \in HV$. Such a 2-cocycle is called a *trivial 2-cocycle* or a *2-coboundary* on HV. Denote by $B^2(HV,\ \mathbb{C})$ the vector space of 2-coboundaries on HV. The quotient space

$$H^2(HV,\ \mathbb{C}) = C^2(HV,\ \mathbb{C})/B^2(HV,\ \mathbb{C})$$

is called the *2-cohomology group* of HV. There exists a one-to-one correspondence between the set of equivalence classes of one-dimensional central extensions of HV by $\mathbb{C}$ and the 2-cohomology group of HV.

Theorem 5.1. *We have* $H^2(HV, \mathbb{C}) = 0$.

Proof It can be easily obtained by [14], Proposition 5.2 and Proposition 5.3. □

Proposition 5.2. *We have*

$$\phi(L_{\alpha,i}, H_{\beta,j}) = 0, \tag{5.1}$$

with $\alpha, \beta \in \Gamma$, $i, j \in \mathbb{Z}_+$.

Proof Without loss of generality, we can always suppose $1 \in \Gamma$. Let $\psi \in C^2(HV, \mathbb{C})$, we define a linear function $f : HV \to \mathbb{C}$ such that $f(L_{\alpha,i})$ and $f(H_{\alpha,i})$ are defined by induction on i as follows

$$f(L_{\alpha,i}) = \begin{cases} \frac{1}{i+1}\psi(L_{0,0}, L_{0,i+1}) & \text{if } \alpha = 0, \\ \frac{1}{\alpha}\Big(\psi(L_{0,0}, L_{\alpha,i}) - if(L_{\alpha,i-1})\Big) & \text{if } \alpha \neq 0. \end{cases} \tag{5.2}$$

$$f(H_{\alpha,i}) = \begin{cases} \psi(L_{0,0}, H_{0,1}) & \text{if } \alpha = i = 0, \\ \frac{1}{i}\psi(L_{0,1}, H_{0,i}) & \text{if } \alpha = 0,\ i \neq 0, \\ \frac{1}{\alpha}\Big(\psi(L_{0,0}, H_{\alpha,i}) - if(H_{\alpha,i-1})\Big) & \text{if } \alpha \neq 0. \end{cases} \tag{5.3}$$

Set $\phi = \psi - \psi_f$, we have

$$\phi(L_{0,0}, H_{\beta,j}) = \psi(L_{0,0}, H_{\beta,j}) - f([L_{0,0}, H_{\beta,j}]) = 0 \ \text{ for } \ \beta \neq 0. \tag{5.4}$$

And we can easily verify that $\phi(L_{0,0}, H_{0,0}) = \phi(L_{0,0}, H_{0,1}) = 0$. Furthermore, for $j \neq 1$ and $j \neq 0$, we obtain

$$\begin{aligned}\phi(L_{0,0}, H_{0,j}) &= \psi(L_{0,0}, H_{0,j}) - jf(H_{0,j-1}) \\ &= \psi(L_{0,0}, H_{0,j}) - \frac{j}{j-1}\phi(L_{0,1}, H_{0,j-1}) \\ &= \psi(L_{0,0}, H_{0,j}) - \frac{1}{j-1}\phi(L_{0,1}, [L_{0,0}, H_{0,j}]) \\ &= 0.\end{aligned}$$

Thus

$$\phi(L_{0,0}, H_{\beta,j}) = 0 \qquad \text{for } \ \beta \in \Gamma, \ j \in \mathbb{Z}_+. \tag{5.5}$$

Now we want to prove $\phi(L_{\alpha,i}, H_{0,0}) = 0$, for any $\alpha \in \Gamma, i \in \mathbb{Z}_+$. If $\alpha \neq 0$, we use induction on i. It holds for $i = 0$. Suppose it also holds for $i-1$, then we get

$$\phi(L_{\alpha,i}, H_{0,0}) = \frac{1}{\alpha}\psi([L_{0,0}, L_{\alpha,i}] - iL_{\alpha,i-1}, H_{0,0}) = 0 \ \text{ for } \quad \alpha \neq 0. \tag{5.6}$$

We can also have

$$\phi(L_{0,i}, H_{0,0}) = \psi(\frac{1}{i+1}[L_{0,0}, L_{0,i+1}], H_{0,0}) = 0 \ \text{ for } \ i \neq 0. \tag{5.7}$$

Therefore, by (5.6) and (5.7), it follows that

$$\phi(L_{\alpha,i}, H_{0,0}) = 0 \qquad \text{for } \ \alpha \in \Gamma, \ i \in \mathbb{Z}_+. \tag{5.8}$$

Similarly, for $\alpha + \beta \neq 0$, we obtain

$$\begin{aligned}\phi(L_{\alpha,0}, H_{\beta,0}) &= \psi(L_{\alpha,0}, H_{\beta,0}) - \beta f(H_{\alpha+\beta,0}) \\ &= \psi(L_{\alpha,0}, H_{\beta,0}) - \frac{\beta}{\alpha+\beta}\psi(L_{0,0}, H_{\alpha+\beta,0}) \\ &= \psi(L_{\alpha,0}, H_{\beta,0}) - \frac{1}{\alpha+\beta}\psi(L_{0,0}, [L_{\alpha,0}, H_{\beta,0}]) \\ &= 0.\end{aligned}$$

By induction on $i + j$, for $\alpha + \beta \neq 0$, we have

$$\begin{aligned}\phi(L_{\alpha,i}, H_{\beta,j}) &= \psi(L_{\alpha,i}, H_{\beta,j}) - \beta f(H_{\alpha+\beta,i+j}) - jf(H_{\alpha+\beta,i+j-1}) \\ &= \psi(L_{\alpha,i}, H_{\beta,j}) - \frac{\beta}{\alpha+\beta}(\psi(L_{0,0}, H_{\alpha+\beta,i+j}) - (i+j)f(H_{\alpha+\beta,i+j-1})) \\ &\quad - \frac{j}{\alpha+\beta}(\psi(L_{0,0}, H_{\alpha+\beta,i+j-1}) - (i+j-1)f(H_{\alpha+\beta,i+j-2})) \\ &= -\frac{1}{\alpha+\beta}(j\phi(L_{\alpha,i}, H_{\beta,j-1}) + i\phi(L_{\alpha,i-1}, H_{\beta,j})) \\ &= 0.\end{aligned}$$

Thus we can conclude that

$$\phi(L_{\alpha,i}, H_{\beta,j}) = \delta_{\alpha+\beta,0}\phi(L_{\alpha,i}, H_{-\alpha,j}) \qquad \text{for } \alpha, \beta \in \Gamma,\ i, j \in \mathbb{Z}_+. \tag{5.9}$$

From the above discussion, we have already known $\phi(L_{0,0}, H_{0,1}) = \phi(L_{0,1}, H_{0,0}) = 0$. Now for the case $i + j > 1$, we have

$$\begin{aligned}
\phi(L_{0,i}, H_{0,j}) &= \psi(L_{0,i}, H_{0,j}) - jf(H_{0,i+j-1}) \\
&= \psi(L_{0,i}, H_{0,j}) - \frac{j}{i+j-1}\psi(L_{0,1}, H_{0,i+j-1}) \\
&= \psi(L_{0,i}, H_{0,j}) - \frac{1}{i+j-1}\psi(L_{0,1}, [L_{0,i}, H_{0,j}]) \\
&= 0.
\end{aligned}$$

So we obtain

$$\phi(L_{0,i}, H_{0,j}) = 0 \qquad \text{for } i, j \in \mathbb{Z}_+. \tag{5.10}$$

In order to complete the proof, by (5.9) and (5.10), we need to compute $\phi(L_{\alpha,i}, H_{-\alpha,j})$ for $\alpha \neq 0$. If $i+j = 1$, we have $\phi(L_{\alpha,0}, H_{-\alpha,1}) = \phi(L_{\alpha,1}, H_{-\alpha,0}) = 0$, for $\alpha \neq 0$. For $i+j > 1$, by induction on $i + j$, it follows that

$$\phi(L_{\alpha,i}, H_{-\alpha,j}) = \psi(L_{\alpha,i}, H_{-\alpha,j}) - \frac{1}{i+j}\psi(L_{0,1}, [L_{\alpha,i}, H_{-\alpha,j}] - jH_{0,i+j-1}) - jf(H_{0,i+j-1}) = 0.$$

Thus

$$\phi(L_{\alpha,i}, H_{-\alpha,j}) = 0 \qquad \text{for } \alpha \neq 0,\ i + j \geq 1. \tag{5.11}$$

Based on the above discussion, we obtain

$$\phi(L_{\alpha,i}, H_{\beta,j}) = \delta_{\alpha+\beta,0}\delta_{i+j,0}\phi(L_{\alpha,0}, H_{-\alpha,0}) \qquad \text{for } \alpha, \beta \in \Gamma,\ i, j \in \mathbb{Z}_+. \tag{5.12}$$

Finally, we consider $\phi(L_{\alpha,0}, H_{-\alpha,0})$. Obviously, we have

$$\begin{aligned}
\phi(L_{\alpha,0}, H_{-\alpha,0}) &= \psi(L_{\alpha,0}, H_{-\alpha,0}) - f([L_{\alpha,0}, H_{-\alpha,0}]) \\
&= \psi(L_{\alpha,0}, H_{-\alpha,0}) + \alpha\psi(L_{0,0}, H_{0,1}) \\
&\triangleq c(\alpha).
\end{aligned}$$

Since $\phi([L_{\alpha,0}, L_{\beta,0}], H_{-\alpha-\beta,0}) + \phi([L_{\beta,0}, H_{-\alpha-\beta,0}], L_{\alpha,0}) + \phi([H_{-\alpha-\beta,0}, L_{\alpha,0}], L_{\beta,0}) = 0$, we get

$$(\beta - \alpha)c(\alpha + \beta) + (\beta + \alpha)c(\alpha) = (\beta + \alpha)c(\beta), \quad \text{for any } \alpha, \beta \in \Gamma. \tag{5.13}$$

In (5.13), replacing α by $\alpha - 1$ and taking $\beta = 1, 2$ respectively, we have

$$\begin{cases}
(2 - \alpha)c(\alpha) + \alpha c(\alpha - 1) = \alpha c(1), \\
(3 - \alpha)c(\alpha + 1) + (\alpha + 1)c(\alpha - 1) = (\alpha + 1)c(2).
\end{cases} \tag{5.14}$$

And by $(1-\alpha)c(\alpha+1)+(\alpha+1)c(\alpha)=(\alpha+1)c(1)$, we obtain

$$c(\alpha)=\frac{c(2)}{2}(\alpha^2-\alpha) \qquad \text{for } \alpha\neq 0. \tag{5.15}$$

As $\phi(L_{0,0},H_{0,0})=0=c(0)$ also satisfies (5.15), we can conclude that (5.15) holds for any $\alpha\in\Gamma$, where $c(2)\in\mathbb{C}$. So $\phi(L_{\alpha,0},H_{-\alpha,0})=b(\alpha^2-\alpha)$, for $b\in\mathbb{C}$.

Finally, using the equality $\phi(L_{0,0},[L_{\alpha,1},H_{-\alpha,0}])=0$, we get $\phi(L_{\alpha,0},H_{-\alpha,0})=0$. □

Proposition 5.3. *We have*

$$\phi(H_{\alpha,i},H_{\beta,j})=0, \tag{5.16}$$

with $\alpha,\beta\in\Gamma$, $i,j\in\mathbb{Z}_+$.

Proof Let $\psi\in C^2(HV,\mathbb{C})$, we define a linear function $f:HV\to\mathbb{C}$ (cf. (5.2) and (5.3)) and set $\phi=\psi-\psi_f$. Then we obtain

$$\phi(H_{0,0},H_{\beta,0})=\psi(H_{0,0},H_{\beta,0})-f([H_{0,0},H_{\beta,0}])=0. \tag{5.17}$$

By induction on j, we have

$$\phi(H_{0,0},H_{\beta,j})=0 \quad \text{for} \quad \beta\in\Gamma,\ j\in\mathbb{Z}_+. \tag{5.18}$$

Furthermore, for $\alpha\neq 0$, we obtain

$$\begin{aligned}\phi(H_{\alpha,0},H_{\beta,0})&=\psi(H_{\alpha,0},H_{\beta,0})-f([H_{\alpha,0},H_{\beta,0}])\\&=\frac{1}{\alpha}\psi([L_{0,0},H_{\alpha,0}],H_{\beta,0})\\&=-\frac{\beta}{\alpha}\phi(H_{\alpha,0},H_{\beta,0}).\end{aligned}$$

Thus $\phi(H_{\alpha,0},H_{\beta,0})=0$ for $\alpha+\beta\neq 0$.

Now considering the case of $\alpha=0$. For $\beta\neq 0$, we also have

$$\begin{aligned}\phi(H_{0,i},H_{\beta,0})&=\psi(H_{0,i},H_{\beta,0})-f([H_{0,i},H_{\beta,0}])\\&=-\frac{(-1)^i i!}{\beta^i}\phi(H_{0,0},H_{\beta,0})\\&=0.\end{aligned}$$

By induction on j, we get

$$\phi(H_{0,i},H_{\beta,j})=0 \quad \text{for} \quad \beta\neq 0. \tag{5.19}$$

Next we want to prove $\phi(H_{\alpha,i},H_{\beta,j})=0$ for $\alpha\neq 0$ and $i+j\geq 1$. Obviously, it holds for $i=0,\ j=1$ and $i=1,\ j=0$. Now assume $\phi(H_{\alpha,i-1},H_{\beta,j})=\phi(H_{\alpha,i},H_{\beta,j-1})=0$. For

$\alpha \neq 0$, considering

$$\begin{aligned}\phi(H_{\alpha,i}, H_{\beta,j}) &= \psi(H_{\alpha,i}, H_{\beta,j}) - f([H_{\alpha,i}, H_{\beta,j}]) \\ &= -\frac{\beta}{\alpha}\phi(H_{\alpha,i}, H_{\beta,j}) - \frac{j}{\alpha}\phi(H_{\alpha,i}, H_{\beta,j-1}) - \frac{i}{\alpha}\phi(H_{\alpha,i-1}, H_{\beta,j}),\end{aligned}$$

it follows that

$$\phi(H_{\alpha,i}, H_{\beta,j}) = 0 \quad \text{for} \quad \alpha + \beta \neq 0, \quad i + j \geq 1. \tag{5.20}$$

Therefore, we obtain

$$\phi(H_{\alpha,i}, H_{\beta,j}) = \delta_{\alpha+\beta,0}\phi(H_{\alpha,i}, H_{-\alpha,j}) \quad \text{for} \quad \alpha \neq 0. \tag{5.21}$$

It is easy to see that $\phi(H_{0,i}, H_{0,j}) = 0$, for $i, j \in \mathbb{Z}_+$. Then we have

$$\phi(H_{\alpha,i}, H_{-\alpha,j}) = -\frac{\alpha}{i-1}\phi(H_{0,j+1}, H_{-\alpha,i}) + \frac{j}{i-1}\phi(H_{0,j}, H_{-\alpha,i}) = 0 \quad \text{for} \quad \alpha \neq 0, \ i \geq 2.$$

Since $\phi(H_{\alpha,0}, H_{-\alpha,1}) = 0$ and $\phi(H_{\alpha,1}, H_{-\alpha,0}) = 0$, we get $\phi(H_{\alpha,i}, H_{-\alpha,j}) = 0$ for $\alpha \neq 0, \ i \geq 1$. It follows that $\phi(H_{\alpha,i}, H_{-\alpha,j}) = \delta_{i+j,0}\phi(H_{\alpha,0}, H_{-\alpha,0})$. Thus we have

$$\phi(H_{\alpha,i}, H_{\beta,j}) = \delta_{\alpha+\beta,0}\delta_{i+j,0}\phi(H_{\alpha,0}, H_{-\alpha,0}) \qquad \text{for} \quad \alpha, \beta \in \Gamma, \ i, j \in \mathbb{Z}_+. \tag{5.22}$$

Now suppose $\phi(H_{\alpha,0}, H_{-\alpha,0}) = g(\alpha)$ and assume $1 \in \Gamma$, we get $\phi(H_{1,0}, H_{-1,0}) = g(1) \in \mathbb{C}$ and $\phi(H_{\alpha,0}, H_{-\alpha,0}) = \psi([L_{\alpha-1,0}, H_{1,0}], H_{-\alpha,0}) = \alpha\psi(H_{1,0}, H_{-1,0}) = \alpha g(1)$, for $g(1) \in \mathbb{C}$.

Finally, using the equality $\phi(L_{0,0}, [H_{\alpha,1}, H_{-\alpha,0}]) = 0$, we get $\phi(H_{\alpha,0}, H_{-\alpha,0}) = 0$. □

References

[1] Arbarello, E., De Concini, C., Kac, V. G., Procesi, C.: Moduli spaces of curves and representation theory, *Comm Math Phys.,* **117**, 1–36(1988)

[2] Billig, Y.: Representations of the twisted Heisenberg-Virasoro algebra at level zero theory, *Canad. Math. Bull.,* **46**, 529–537(2003)

[3] Chen, Q., Han, J., Su, Y.: Structures of not-finitely graded Lie algebras related to generalized Virasoro algebras, *Comm. Algebra.,* **43**, 3033–3049(2015)

[4] Dokovic, D., Zhao, K.: Derivations, isomorphisms, and second cohomology of generalized Witt algebras. *Tran. Ams. Math. Soc.,* **350**, 643–664(1998)

[5] Farnsteiner, R.: Derivations and central extensions of finitely generated graded Lie algebras, *J. Algebra.,* **118**, 33–45(1988)

[6] Kac, V.G.: *Infinite-dimensional Lie algebras*, (3rd edn), Cambridge University Press, Cambridge, 1990

[7] Kac, V. G., Peterson, D. H.: Spin and representation of infinite dimensional Lie algebras and groups. *Proc. Nat. Acad. Sci. U.S.A.,* 3308–3312(1981)

[8] Liu, D., Wu, Y., Zhu, L.: Whittaker modules for the twisted Heisenberg-Virasoro algebras, *J. Math.Phys.,* **51**, 1–12(2010)

[9] Liu, D., Zhu, L.: Generalized Heisenberg-Virasoro algebras, *Front. Math.China.,* **4**, 297–310(2009)

[10] Shen, R., Jiang, C.: Derivation algebra and automorphism group of the twisted Heisenberg-Virasoro algebra, *Comm. Algebra.,* **34**, 2547–2558(2006)

[11] Su, Y.: 2-Cocycles on the Lie algebras of generalized differantial operators, *Comm. Algebra.,* **30**, 763–782(2002)

[12] Su, Y.: Derivations of generalized Weyl algebras, *Science in China A.,* **46**, 346–354(2003)

[13] Su, Y., Xu, X., Zhang, H.: Derivation-simple algebras and the structures of Lie algebras of Witt type, *J. Algebra.,* **233**, 642–662(2000)

[14] Su, Y., Zhao, K.: Second cohomology group of generalized Witt type Lie algebras and certain representations, *Comm. Algebra.,* **30** , 3285–3309(2002)

[15] Su, Y., Zhao, K.: Isomorphism classes and automorphism groups of algebras of Weyl type, *Science in China A.,* **32**, 954–963(2002)

[16] Su, Y., Zhou, J.: Structure of the Lie algebras related to those of Block, *Comm. Algebra.,* **30**, 3205–3226(2002)

[17] Wu, H., Wang, S., Yue, X.: Structures of generalized loop Virasoro algebras, *Comm. Algebra.,* **42**, 1545–1558(2014)